



ESMO 2026

Zoci (ZL-1310): 有望成为小细胞肺癌和神经内分泌癌治疗的基石药物

2026年10月

NASDAQ:ZLAB | HKEX:9688



© 2026. Zai Lab. All Rights Reserved.

前瞻性声明

本演示材料包含前瞻性陈述，包括与以下方面相关的陈述：我们的策略和计划；我们的业务、商业化产品及管线项目的潜力和预期；特别是 zoci（ZL-1310），我们的目标、目的和重点事项，以及我们基于增长战略的预期（包括我们对商业化产品和上市以及 zoci 等临床阶段产品的预期）；我们项目的峰值销售潜力；资本配置与投资策略；临床开发计划和相关临床研究，包括临床研究的计划启动、设计和数量，以及我们与合作伙伴的联合治疗方案的计划；临床研究数据、数据解读和发布的预期时间与结果；注册相关的讨论、提交、申请、获批及时间线，包括新药临床试验申请（IND）的预期时间、我们首个生物制品许可申请（BLA）的潜在提交以及潜在的全球批准；我们及我们合作伙伴的产品和候选产品的潜在裨益、安全性和有效性；投资、合作及业务拓展活动的预期收益和潜力；我们商业化产品和候选产品的潜在市场机会及预估的可及市场，包括预估患者人群；以及我们未来的财务和经营业绩。除对过往事实的陈述外，本演示材料中包含的所有陈述均为前瞻性陈述，可以通过诸如“旨在”、“预计”、“认为”、“有可能”、“估计”、“预期”、“预测”、“目标”、“打算”、“可能”、“计划”、“可能的”、“潜在”、“将”、“将会”等词汇及其他类似表述加以识别。该等陈述构成美国联邦证券法所定义的前瞻性陈述。前瞻性陈述并非对未来表现的担保或保证，因为预测未来结果存在固有困难。

前瞻性陈述基于我们截至本演示材料发布之日的预期和假设，并且受到固有的不确定性、风险以及可能与前瞻性陈述所预期的情况存在重大差异的情势变更的影响。我们可能无法实际实现在我们的前瞻性陈述中所披露的计划、意图、预期或预测，您不应过分依赖该等前瞻性陈述。实际结果可能受各多种重要因素的影响而与前瞻性陈述所示存在重大差异，这些因素包括但不限于：(1) 我们成功商业化自身已获批产品并从中产生收入的能力；(2) 我们为自身的运营和业务活动获取资金的能力；(3) 我们候选产品临床及临床前开发的结果，包括初步或中期数据可能无法预示最终或后期结果以及跨研究比较可能缺乏意义的风险；(4) 相关监管机构对我们候选产品作出审批决定的内容和时间；(5) 来自其他制药和生物技术公司的产品及候选产品的潜在竞争；(6) 我们所依赖的第三方（如我们的许可方和合作伙伴）可能无法成功履行其合同义务或按预期期限完成工作；(7) 与在中国营商有关的风险；以及 (8) 我们向美国证券交易委员会（SEC）提交的最新年报和季报及其他报告中指出的其他因素。我们预计后续事件和发展将导致我们的预期和假设发生变化，但除法律要求之外，无论出于新信息、未来事件或其他原因，我们均无义务更新或修订任何前瞻性陈述。

市场与行业数据

除另有说明外，本演示材料中包含的市场与行业数据（包括预估的可及市场、患者人群和潜在市场机会）均基于管理层的估计和研究，以及第三方开展的行业与综合出版物、研究、调查和研究报告。我们并未对这些信息的准确性或完整性进行独立核实。管理层的估计源于公开可得的信息、其对所处行业的了解，以及基于此类信息和了解所作的假设。这些数据涉及若干假设和局限性，且由于多种因素，我们所处的行业存在高度的不确定性和风险。

商标

本演示材料中出现的商标和注册商标均为其各自所有者的财产。

今日讲者与议程



Rafael Amado, 医学博士

再鼎医药总裁，全球研发负责人



Pedro Rocha, 医学博士，哲学博士，理学硕士

西班牙巴塞罗那 Vall d'Hebron 大学医院肿瘤内科胸部肿瘤科

介绍

Rafael Amado, 医学博士

Zoci: 2026 ESMO 亮点 – 一线小细胞肺癌

Pedro Rocha, 医学博士，哲学博士，理学硕士

总结和问答

Rafael Amado, 医学博士

Pedro Rocha, 医学博士，哲学博士，理学硕士

Pedro Rocha, 医学博士, 哲学博士, 理学硕士

- 接受以下公司的酬金：雅培、安进、阿斯利康、第一三共、礼来、赛默飞。
- 担任以下公司的咨询或顾问角色：第一三共、IGES Pharma。
- 接受以下公司提供的差旅和住宿费用：阿斯利康、百时美施贵宝、第一三共、协和麒麟、默沙东、PharmaMar、罗氏、赛默飞。

持续拓展全球肿瘤与免疫管线，数据读出保持稳定节奏

肿瘤

免疫


项目（靶点）		适应证	IND支持阶段	I期	概念验证	注册性研究/III期	当前状态/下一次更新
Zoci (DLL3 ADC)	Zoci	2L+ 小细胞肺癌				●	数据（中期分析）：2027年下半年
	Zoci + PD-(L)1	1L 小细胞肺癌			●		数据已于ESMO大会公布；III 期研究计划中
	Zoci + tarlatamab (TCE) ¹ ± PD-L1	1L+ 小细胞肺癌		●			Ib 期研究正在进行中
	Zoci + obixtamig (TCE) ² ± PD-L1	小细胞肺癌及肺外神经内分泌癌		●			Ib/II 期研究将于2026年下半年启动
	Zoci	肺外神经内分泌癌				●	数据（注册性研究）：2027年上半年
ZL-6201 (LRRRC15 ADC)	ZL-6201	肉瘤及其他肿瘤			●		数据（概念验证）：2027年上半年
ZL-1222 (PD-1/IL-12)	ZL-1222	实体瘤	●				I 期研究将于2027年启动
ZL-1311 (MUC17/CD3 TCE)	ZL-1311	实体瘤	●				I 期研究将于2027年上半年启动
ZL-6202 (未披露)	ZL-6202	实体瘤	候选药物遴选				
ZL-1223 (未披露)	ZL-1223	实体瘤	候选药物遴选				
ZL-1503 (IL-13/IL-31Rα)	ZL-1503	中重度特应性皮炎			●		数据（健康志愿者SAD）：2026年第四季度 数据（特应性皮炎患者MAD）：2027年上半年
ZL-1505 (未披露)	ZL-1505	免疫性疾病	候选药物遴选				

注：SAD：单次递增剂量；MAD：多次递增剂量。（1）安进负责申办并主导全球Ib/II期研究；再鼎医药提供研究药物。（2）勃林格殷格翰负责申办并主导全球Ib/II期研究；再鼎医药提供研究药物。

Zoci: 兼具强效抗肿瘤活性与良好耐受性的新一代ADC

Zoci基于TMALIN[®]平台构建，实现强效、精准的肿瘤杀伤¹

ZL-1310的结构与作用机制



A: 人源化抗DLL3 IgG1单克隆抗体

B: 可裂解的三肽连接子

C: 喜树碱衍生物载荷C24

Zocilurtatug pelitecan（简称zoci）是一款靶向DLL3的新型ADC。DLL3是一种神经内分泌肿瘤特异性抗原，在小细胞肺癌中呈高表达²⁻⁴，并已被验证为具有治疗价值的靶点。

关键特性

DAR=8

向靶细胞高效递送载荷

肿瘤微环境特异性释放

通过肿瘤微环境可激活连接子减少全身毒性，提升耐受性


强效旁观者杀伤效应

Topo-1抑制剂载荷C24可增强抗肿瘤作用，包括对异质性肿瘤的杀伤


五大特性使zoci有望成为小细胞肺癌和神经内分泌癌治疗的基石ADC

差异化的临床特征、正在开展的全球III期研究，以及覆盖小细胞肺癌和神经内分泌癌的广泛开发计划

01




高缓解率



02




持久缓解



03




潜在同类最佳的
安全性



04




强劲的颅内
抗肿瘤活性



05



全球领先的
临床开发进展



强劲疗效 • 良好耐受性 • 颅内抗肿瘤活性 • 全球开发

DLL3 ADC

Zoci (ZL-1310)

ESMO 2026 Data Update

Pedro Rocha, MD, PhD, MSc
Vall d'Hebron University Hospital

ZL-1310 (DLL3 ADC) plus atezolizumab $\pm$ carboplatin in Extensive-stage Small Cell Lung Cancer

Zhao Jun, MD

Key Laboratory of Carcinogenesis and Translational Research
(Ministry of Education/Beijing), Peking University Cancer Hospital
and Institute, Beijing, China



Jun Zhao¹, Valentina Gambardella², Wenxiu Yao³, Zhen Wang⁴, Pedro Rocha⁵,
Jie Hu⁶, María Eugenia Olmedo García⁷, Qiming Wang⁸, David Vicente Baz⁹,
Luis Paz-Ares¹⁰, Mariam Alexander¹¹, Anne C. Chiang¹², Manuel Cobo Dols¹³,
Yingying Du¹⁴, Grace K. Dy¹⁵, Martin Gutierrez¹⁶, Alexander Spira¹⁷, Liza
Villaruz¹⁸, Renke Zhou¹⁹, Yi-Long Wu³

¹Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Peking University Cancer Hospital and Institute, Beijing, China; ²Hospital Clínico Universitario de Valencia, INCLIVA, Valencia, Spain; ³Sichuan Cancer Hospital, Chengdu, China; ⁴Guangdong Lung Cancer Institute, Guangdong Provincial People's Hospital (Guangdong Academy of Medical Sciences), Southern Medical University, Guangzhou, China; ⁵Instituto de Investigacion Oncologica Vall d'Hebron (VHIO), Barcelona, Spain; ⁶Shanghai Geriatric Medical Center, Zhongshan Hospital, Fudan University, Shanghai, China; ⁷Hospital Universitario Ramón y Cajal, Madrid, Spain; ⁸Henan Cancer Hospital, Zhengzhou, China; ⁹Hospital Universitario Virgen Macarena, Sevilla, Spain; ¹⁰Hospital Universitario 12 de Octubre, Madrid, Spain; ¹¹Medical University of South Carolina, Charleston, SC, USA; ¹²Yale Cancer Center, Yale University School of Medicine, New Haven, CT, USA; ¹³Hospital Regional Universitario de Málaga, Málaga, Spain; ¹⁴The First Affiliated Hospital of Anhui Medical University, Hefei, China; ¹⁵Roswell Park Comprehensive Cancer Center, Buffalo, NY, USA; ¹⁶Hackensack University Medical Center, Hackensack, NJ, USA; ¹⁷NEXT Oncology Virginia, Fairfax, VA, USA; ¹⁸UPMC Hillman Cancer Center, Pittsburgh, PA, USA; ¹⁹Zai Lab (US) LLC, Cambridge, USA

Study Overview

- Phase I, open-label, dose-escalation and expansion study of ZL-1310 as monotherapy and in combination with atezolizumab, with (triplet) or without (doublet) carboplatin in ES-SCLC (NCT06179069).
- Efficacy and safety reported for 1L patients who had the opportunity of at least 12-week follow-up with data cutoff date of Sept 9, 2026.

ZL-1310-001 Study Schema (ZL-1310 plus atezolizumab ± carboplatin)


• Patient population

• Doublet:

- **Dose escalation:** relapsed / refractory ES-SCLC ≤2 prior lines of therapy.
- **Backfill:** Newly diagnosed ES-SCLC (1L) or ES-SCLC completed standard 1L induction therapy and had not progressed (1L maintenance).
- **Expansion:** Newly diagnosed ES-SCLC (1L)

• Triplet: Newly diagnosed ES-SCLC (1L)

- Asymptomatic brain metastasis (treated or untreated) allowed
- Prior DLL3-targeted therapy allowed
- ECOG Performance Status 0-1



Baseline Characteristics for 1L Patients

ZL-1310 + atezolizumab doublet:

- At ZL-1310 1.6 mg/kg, 33 patients were enrolled; the first 15 in the back fill cohort had the opportunity for ≥ 6 months of follow-up, with a median follow-up of 10.0 months.
- At ZL-1310 1.2 mg/kg, 10 patients were enrolled; all had ≥ 6 months follow-up, with a median follow-up of 12.6 months.

ZL-1310 + atezolizumab + carboplatin triplet:

- In the dose-escalation triplet cohorts, 13 patients were enrolled, with median follow-up of 11.8 months.

Demographics and Baseline Characteristics for 1L Patients

ZL-1310 + atezolizumab doublet				ZL-1310 + atezolizumab + carboplatin triplet			
	Dose Level		Total	Dose Level			Total
	ZL-1310			ZL-1310			
				Carbo AUC5	Carbo AUC4		
	1.2 mg/kg (n=10)	1.6 mg/kg (n=33)	Total (n=43)	0.8 mg/kg (n=4)	0.8 mg/kg (n=3)	1.2 mg/kg (n=6)	Total (n=13)
Male, n (%)	8 (80.0)	26 (78.8)	34 (79.1)	3 (75.0)	3 (100)	4 (66.7)	10 (76.9)
Age, median (range), years	62 (51, 75)	63 (46, 75)	63 (46, 75)	61 (58, 66)	69 (66, 69)	68 (59, 75)	66 (58, 75)
ECOG PS = 1, n (%)	7 (70.0)	25 (75.8)	32 (74.4)	3 (75.0)	3 (100)	5 (83.3)	11 (84.6)
Race, n (%)							
Asian	10 (100)	30 (90.9)	40 (93.0)	3 (75.0)	2 (66.7)	2 (33.3)	7 (53.8)
White	0	3 (9.1)	3 (7.0)	0	1 (33.3)	4 (66.7)	5 (38.5)
Brain Metastasis-Yes, n (%)	2 (20.0)	6 (18.2)	8 (18.6)	1 (25.0)	0	2 (33.3)	3 (23.1)
Stage IV disease at initial diagnosis, n (%)	8 (80.0)	25 (75.8)	33 (76.7)	3 (75.0)	2 (66.7)	5 (83.3)	10 (76.9)

Results: Target Tumor Reduction of Doublet in 1L

- High overall response rates were observed in patients receiving ZL-1310 + atezolizumab as 1L therapy (ORR for all dose level: 81.4%).

Target Tumor Reduction per RECIST v1.1 by Investigator Assessment




Table 2. Best Overall Response per RECIST v1.1 by Investigator Assessment


Tumor response	ZL-1310 1.2 mg/kg + Atezo (N=10)	ZL-1310 1.6 mg/kg + Atezo (N=33)
Best overall response, n (%)		
CR	0	0
PR	8 (80.0)	27 (81.8)
SD	2 (20.0)	2 (6.1)
PD	0	2 (6.1)
cORR, % (95% CI)	80.0 (44.4, 97.5)	81.8 (64.5, 93.0)

* Patients without target lesions at baseline or evaluable post-baseline tumor assessments were not shown. As of the DCO, two patients reached EOT without disease progression before an evaluable post-baseline assessment was obtained.

Results: Durability of Tumor Reduction with Doublet

- Responses occurred rapidly: 34/35 (97.1%) responders achieving their response at the first post-baseline tumor assessment.
- The responses were durable.
 - In the 10 patients receiving ZL-1310 1.2 mg/kg + atezolizumab as 1L, mDoR was 8.3 (4.0, NA) months.
 - In the first 15 patients receiving ZL-1310 1.6 mg/kg + atezolizumab as 1L, mDoR is not reached after a median follow-up of 10.0 months.

Durable Tumor Reduction Observed in Doublet Cohorts



All 33 patients on 1.6 mg/kg doublet

81.9%

6-month PFS rate

75.8%

Of patients still on treatment as of DCO (25/33)

In the first 15 patients of 1.6mg/kg doublet in 1L: (median follow-up: 10 mo)

66.7%


Have not yet experienced disease progression or death at DCO

NR

Median PFS

Results: Target Tumor Reduction of Triplet in 1L

- Patients receiving ZL-1310 in combination with atezolizumab and carboplatin as 1L therapy demonstrated a high overall response rate (ORR) of 92.3%.




* Patients without target lesions at baseline or evaluable post-baseline tumor assessments were not shown. As of the DCO, one patient in triplet cohort reached EOT without disease progression before an evaluable post-baseline assessment.

Results: Durability of Tumor Reduction with Triplet

- Patients receiving ZL-1310 in combination with atezolizumab and carboplatin as 1L therapy demonstrated a high overall response rate (ORR) of 92.3%.
- mPFS is 8.4 months (95% CI 3.7, NA) and mDoR is 7.7 months (2.6, NA).

Durable Tumor Reduction Observed in Triplet Cohort



Results: Safety of Doublet as 1L Therapy

ZL-1310 plus atezolizumab is well tolerated.

- No DLTs were observed in the dose escalation phase.
- The majority of AEs were Grade 1 or 2, and non-serious.
- Grade ≥ 3 AEs related to ZL-1310 occurred in 7/43 (16.3%) of patients, and SAEs related to ZL-1310 occurred in 5/43 (11.6%) of patients.
- AE rate related to ZL-1310 that led to dose reduction (4.7%, 2/43) or discontinuation (1/43, 2.3%) was low.
- No treatment-related deaths were reported.

Summary of Adverse Events - Doublet Combination Safety Analysis Set

AEs, n (%)	ZL-1310 1.2 mg/kg + Atezolizumab (N=10)	ZL-1310 1.6 mg/kg + Atezolizumab (N=33)	Total (N=43)
Any	10 (100)	32 (97.0)	42 (97.7)
Related to ZL-1310	10 (100)	31 (93.9)	41 (95.3)
Serious	4 (40.0)	6 (18.2)	10 (23.3)
Related to ZL-1310	2 (20.0)	3 (9.1)	5 (11.6)
Grade ≥ 3	5 (50.0)	7 (21.2)	12 (27.9)
Related to ZL-1310	4 (40.0)	3 (9.1)	7 (16.3)
Leading to ZL-1310 Drug Interruption ⁽¹⁾	5 (50.0)	9 (27.3)	14 (32.6)
Related to ZL-1310	4 (40.0)	5 (15.2)	9 (20.9)
Leading to ZL-1310 Dose Reduction	0	2 (6.1)	2 (4.7)
Related to ZL-1310	0	2 (6.1)	2 (4.7)
Leading to ZL-1310 Drug Discontinuation	1 (10.0)	0	1 (2.3)
Related to ZL-1310	1 (10.0)	0	1 (2.3)

* As dose expansion of ZL-1310 + atezolizumab was ongoing at the data cutoff, safety data remain preliminary and subject to ongoing data cleaning. (1) Drug interruption includes infusion interruption or treatment delay compared with the scheduled date due to AE.

Results: Safety of Triplet as 1L Therapy

Compared with the doublet, the triplet was associated with increased hematologic toxicity, more frequent Grade ≥ 3 AEs, and a higher rate of dose interruptions. Although MTD was not met, the ZL-1310 dose was not escalated above 1.2 mg/kg.

- A single DLT, Grade 4 thrombocytopenia, was observed in ZL-1310 0.8 mg/kg + Atezolizumab + Carboplatin AUC 5. No DLTs occurred in the ZL-1310 + Atezolizumab + Carboplatin AUC4 cohorts.
- Grade ≥ 3 AEs related to ZL-1310 occurred in 7/13 (53.8%) of participants, and SAEs related to ZL-1310 occurred in 3/13 (23.1%).
- One patient from ZL-1310 1.2 mg/kg + Atezolizumab + Carboplatin AUC 4 cohort had over-dose of Carboplatin and treatment-related death due to a G5 Sepsis.

Summary of Adverse Events - Triplet Combination Safety Analysis Set

AEs, n (%)	ZL-1310 0.8 mg/kg + Atezolizumab + Carboplatin AUC 5 (N=4)	ZL-1310 0.8 mg/kg + Atezolizumab + Carboplatin AUC 4 (N=3)	ZL-1310 1.2 mg/kg + Atezolizumab + Carboplatin AUC 4 (N=6)	Total (N=13)
Any	4 (100)	3 (100)	6 (100)	13 (100)
Related to ZL-1310	3 (75.0)	2 (66.7)	5 (83.3)	10 (76.9)
Serious	2 (50.0)	0	5 (83.3)	7 (53.8)
Related to ZL-1310	1 (25.0)	0	2 (33.3)	3 (23.1)
Grade ≥ 3	3 (75.0)	1 (33.3)	6 (100)	10 (76.9)
Related to ZL-1310	2 (50.0)	1 (33.3)	4 (66.7)	7 (53.8)
Leading to ZL-1310 Drug Interruption ⁽¹⁾	4 (100)	3 (100)	4 (66.7)	11 (84.6)
Related to ZL-1310	3 (75.0)	2 (66.7)	4 (66.7)	9 (69.2)
Leading to ZL-1310 Dose Reduction	0	0	1 (16.7)	1 (7.7)
Related to ZL-1310	0	0	1 (16.7)	1 (7.7)
Leading to ZL-1310 Drug Discontinuation	0	0	1 (16.7)	1 (7.7)
Related to ZL-1310	0	0	1 (16.7)	1 (7.7)

(1) Drug interruption includes infusion interruption or treatment delay compared with the scheduled date due to AE.

Common AEs related to ZL-1310 and ILD/pneumonitis


Common ZL-1310 Related AE:

- Doublet: The most common AEs related to ZL-1310 of all grades (≥20%) were anaemia (48.8%), nausea (39.5%), leukopenia* (34.9%), decreased appetite (30.2%), ALT increased, neutropenia* (27.9% each), hypertriglyceridaemia and lymphopenia* (23.3% each)
- Triplet: The most common AEs related to ZL-1310 of all grades (≥20%) were anaemia and neutropenia* (61.5% each), thrombocytopenia* (53.8%), nausea (46.2%), ALT increased, decreased appetite, and leukopenia* (38.5% each), AST increased, constipation, malaise, and vomiting (30.8% each), and alopecia, hyperuricaemia, and hypoaesthesia (23.1% each).

ILD / pneumonitis

- ILD / pneumonitis was low in both regimens: 2/43 (4.7%) in doublet, and 2/13 (15.4%) in triplet, all Grade 1. Only one was reported so far in patients receiving ZL-1310 1.6 mg/kg and atezolizumab as 1L treatment (1/33, 3.0%).

Common TRAEs related to ZL-1310 by grade (≥20%)



Leukopenia* includes AEs coded as Leukopenia and White blood cell count decreased. Neutropenia* includes AEs coded as Neutropenia and Neutrophil count decreased. Lymphopenia* includes AEs coded as Lymphocyte count decreased. Thrombocytopenia* includes AEs coded as Platelet count decreased and Thrombocytopenia.

Conclusion

- ZL-1310 1.6 mg/kg and atezolizumab doublet combination demonstrated high response rate (81.8%) and durable response as 1L treatment in ES-SCLC (median PFS for the first 15 patients who had 10-month follow-up not yet obtained).
- ZL-1310 1.6 mg/kg and atezolizumab doublet combination is well tolerated. The overall AE rate, Grade 3 and above AE rate, SAE rate with this regimen is not higher than that of ZL-1310 1.6 mg/kg monotherapy as previously reported. Only a G1 ILD / pneumonitis was reported so far in patients receiving ZL-1310 1.6 mg/kg and atezolizumab as 1L treatment (1/33, 3.0%).
- ZL-1310 1.6 mg/kg and atezolizumab doublet combination has more favorable risk / benefit profile compared to ZL-1310 + atezolizumab + carboplatin triplet regimen.
- A Phase III randomized 1L study will be started to further evaluate the combination regimen in ES-SCLC (NCT07796100).

Summary

Rafael Amado, MD

President, Head of Global R&D

一线小细胞肺癌治疗格局持续演进，zoci战略优势地位凸显



Zoci + DLL3 TCE：两项全球研究探索一线三联方案用于小细胞肺癌和神经内分泌癌



- 安进和勃林格殷格翰将分别负责申办并主导各自的全球Ib/II期研究；再鼎医药将提供研究药物
- 再鼎医药将继续探索其他联合治疗方案

推进zoci在多个适应证和治疗阶段的开发


适应证	治疗线次	治疗方案	研究阶段	开发状态
广泛期小细胞肺癌（ES-SCLC）	2L+	Zoci 单药	III 期	III期研究正在进行中
	1L	Zoci + PD-L1	I 期	I期研究正在进行中；III期研究计划中
	1L+	Zoci + TCE ± PD-L1	Ib 期	Ib期研究正在进行中
肺外神经内分泌癌（epNEC）	2L+	Zoci 单药	I/II 期	注册性队列已完成全部患者入组
	1L+	Zoci + TCE ± PD-L1	I 期	I期研究启动中

其他计划探索的适应证：局限期小细胞肺癌（LS-SCLC）、脑瘤、肺大细胞神经内分泌癌（LCNEC）

Zoci与ZL-1503：更多拓展机遇

数据读出与监管注册申报稳步推进，未来拓展机遇持续显现

当前



Note: SCLC = small cell lung cancer; epNEC = extrapulmonary neuroendocrine carcinomas

Q&A

APPENDIX

Abbreviations

BLA	Biologics License Application	LS-SCLC	Limited-stage small cell lung cancer
CAF	Cancer-associated fibroblast	mAb	Monoclonal antibody
CR	Complete response	MAD	Multiple ascending dose
DAR	Drug-to-antibody ratio	MOA	Mechanism of action
DLL3	Delta-like ligand 3	NEC	Neuroendocrine carcinoma
DoR	Duration of response	OMG	Ocular myasthenia gravis
EP	Etoposide and platinum	ORR	Objective response rate
epNEC	Extrapulmonary neuroendocrine carcinoma	pMN	Primary membranous nephropathy
ES-SCLC	Extensive-stage small cell lung cancer	PS	Performance status
GC	Gastric cancer	Q3W	Once every 3 weeks
GEJ	Gastroesophageal junction	SAD	Single ascending dose
Ig	Immunoglobulin	SCLC	Small cell lung cancer
IgAN	Immunoglobulin A nephropathy	TME	Tumor microenvironment
ILD	Interstitial lung disease	TRAE	Treatment-related adverse event
IO	Immuno-oncology		
LCNEC	Large cell neuroendocrine carcinoma		
LRRC15	Leucine-rich repeat-containing protein 15		